

University of Groningen

Nutritional and reproductive strategies in a chemosymbiotic bivalve living in a tropical intertidal seagrass bed

van der Geest, Matthijs; Sall, Amadou Abderahmane; Ely, Sidi Ould; Nauta, Reindert W.; Gils, Jan A. van; Piersma, Theunis

Published in:
Marine Ecology Progress Series

DOI:
[10.3354/meps10702](https://doi.org/10.3354/meps10702)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Early version, also known as pre-print

Publication date:
2014

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van der Geest, M., Sall, A. A., Ely, S. O., Nauta, R. W., Gils, J. A. V., & Piersma, T. (2014). Nutritional and reproductive strategies in a chemosymbiotic bivalve living in a tropical intertidal seagrass bed. *Marine Ecology Progress Series*, 501, 113-126. <https://doi.org/10.3354/meps10702>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Nutritional and reproductive strategies in a chemosymbiotic bivalve living in a tropical intertidal seagrass bed

Matthijs van der Geest^{1,*}, Amadou Abderahmane Sall², Sidi Ould Ely³, Reindert W. Nauta¹, Jan. A. van Gils¹, Theunis Piersma^{1,4}

¹NIOZ Royal Netherlands Institute for Sea Research, PO Box 59, 1790 AB Den Burg, Texel, The Netherlands

²Mauritanian Institute for Oceanographic Research and Fisheries (IMROP), BP 22, Nouadhibou, Mauritania

³Parc National du Banc d'Arguin, BP 5355, Nouakchott, Mauritania

⁴Chair in Global Flyway Ecology, Animal Ecology Group, Centre for Ecological and Evolutionary Studies (CEES), University of Groningen, PO Box 11103, 9700 CC Groningen, The Netherlands

*Corresponding author: matt.van.der.geest@gmail.com

Marine Ecology Progress Series 501: 113–126 (2014)

Supplement. This supplement contains a short habitat characterization of the study site at Banc d'Arguin, Mauritania (Text S1) and several figures. These figures include information about the size of sexual differentiation and maturity in *Loripes lucinalis* (Fig. S1), the size at first reproduction in *L. lucinalis* (Fig. S2), and the relationship between environmental factors (i.e. air temperature and precipitation), feeding mode, and variables describing the body mass and reproductive cycle of *L. lucinalis* (Fig. S3).

Text S1: Sediment characteristics of the study site

To characterize the habitat at our study site, we took 4 replicate 8 cm deep sediment cores (2 cm diam.) on 7 October 2010 for analysis of median grain size (MGS) and percentage of silt/clay (particles <63 µm). Our study site can be characterized by sediment with MGS of 51.4 µm (range 43.0 to 56.3) and with a percentage of silt/clay of 62.5% (V/V; range 57.2 to 72.7). The reported sediment characteristics at our study site are well within the range of earlier observed values for seagrass-covered intertidal sediments at Banc d'Arguin, Mauritania (van der Geest et al. 2011).




Fig. S1. *Loripes lucinalis*. Percentage of females, males and undifferentiated *L. lucinalis* per size-class (mm). Numbers above columns indicate samples sizes




Fig. S2. *Loripes lucinalis*. Percentage of gonads in 'spent' stage per size-class (mm) for mature *L. lucinalis* (≥ 7 mm in shell height, see Fig. S1) collected in February and July 2010 (i.e. the months in which the 2 major spawning events occurred; see Fig. 4 in the main text). Numbers above columns indicate samples sizes




Fig. S3. *Loripes lucinalis*. Matrix of scatterplots (lower panels) and Pearson's coefficients (upper panels) of correlations between body mass index (BMI, mg cm^{-3}), somatic mass index (SMI, mg cm^{-3}), gonadal mass index (GMI, mg cm^{-3}), percentage of mature *L. lucinalis* individuals classified as 'spent' (Spent), relative heterotrophic contribution to the diet of mature individuals (Diet), air temperature (T, $^{\circ}\text{C}$), precipitation (P, mm), and between month changes in temperature (ΔT , $^{\circ}\text{C}$), and precipitation (ΔP , mm), respectively. Monthly mean values ($n = 15$) were used for all variables (note that for the plots showing between-month changes in abiotic, factors $n = 14$). GMI data were square-root-transformed, and data expressed in percentages (Spent and Diet) were logit-transformed to obtain normality. Significant correlations are indicated with a reduced major axis regression line through the data. Bold values represent significant correlations ($p < 0.05$)